

FDI policy statement: Minimal Intervention in the management of dental caries

Adopted by the FDI General Assembly: 1 October 2002 – Vienna



The FDI World Dental Federation supports the principles of minimal intervention dentistry in the management of dental caries.

FDI

世界牙科联盟支持在牙科龋患处理中使用最小干预牙科的原则。

La Federación Dental Internacional (FDI), apoya los principios de la mínima intervención en odontología para el manejo de caries dentales.

These principles are:

1. Modification of the oral flora

Dental caries is an infectious disease, and the primary focus should therefore be on control of the infection, plaque control and reduced carbohydrate intake.

2. Patient education

The aetiology of dental caries should be explained to the patient, together with the means of prevention through dietary and oral hygiene measures.

3. Remineralisation of non-cavitated lesions of enamel and dentine

Saliva plays a critical role in the demineralisation / remineralisation cycle, and its quantity and quality should therefore be assessed. There is strong evidence that 'white spot' lesions of enamel and non-cavitated lesions of dentine can be arrested or reversed. Such lesions should therefore be managed initially by remineralisation techniques. The extent of the lesion should be objectively recorded such that any progression can be identified at recall.

4. Minimal operative intervention of cavitated lesions

An operative ('surgical') approach should only be used when specifically indicated, e.g. when cavitation is such that the lesion cannot be arrested, or when there are aesthetic or functional requirements. Operative intervention should focus on the preservation of natural tooth structure and be limited to the removal of friable enamel and infected dentine. This can be done with hand, rotary, sonic, ultrasonic, air abrasive or laser instruments, depending on the circumstances. Each prepared cavity is therefore unique, and is primarily dependent on the extent of infected dentine rather than on a predetermined cavity design. Preparation of minimal cavities enables their restoration with adhesive materials such as glass-ionomer cement and/or resin composite. Some studies suggest that glass-ionomer cement may aid in the remineralisation of demineralised, firm, non-infected dentine; however, further clinical studies are needed.

5. Repair of defective restorations

Removal of restorations results in an inevitable increase in cavity size as a consequence of removal of sound tooth structure. Depending on the clinical judgment of the dentist, repair could be considered as an alternative to replacement in some circumstances.

Reference

Tyas M J, Anusavice K J, Frencken J E, Mount G J. Minimal Intervention Dentistry - A Review. FDI Commission Project 1-97. Int Dent J 2000; 50: 1-12.

W:\Exec
Office\Statements\Website
Statements\ENGLISH\Minimal Intervention in
the Management of Dental Caries.doc

Version dated: 1 October 2002

http://www.fdiworldental.org/federation/assets/statements/ENGLISH/Caries/Minimal_intervention_caries.pdf